



An indecomposable PD_3 -complex : II

JONATHAN A. HILLMAN

Abstract We show that there are two homotopy types of PD_3 -complexes with fundamental group $S_3 *_{\mathbb{Z}/2\mathbb{Z}} S_3$, and give explicit constructions for each, which differ only in the attachment of the top cell.

AMS Classification 57P10; 55M05

Keywords Indecomposable, Poincaré duality, PD_3 -complex

In [3] we showed that $\pi = S_3 *_{\mathbb{Z}/2\mathbb{Z}} S_3$ satisfies the criterion of [5] and thus is the fundamental group of a PD_3 -complex. As π has infinitely many ends but is indecomposable, this illustrates a divergence from the known properties of 3-manifolds, and provides a counter-example to an old question of Wall [6]. In particular, the Sphere Theorem does not extend to all PD_3 -complexes.

Here we shall give an explicit description of a finite PD_3 -complex Y realizing this group. The construction is modelled on a similar construction for a PD_3 -complex X with fundamental group S_3 . In each case the cellular chain complex of the universal cover has the striking property that it is self-dual. In §2 we show a PD_3 -complex with fundamental group π must be orientable, and we use Turaev's work to show there are two homotopy types of such PD_3 -complexes. The 2-fold cover of Y is homotopy equivalent to $L(3, 1) \# L(3, 1)$, while a simple modification of our construction (suggested by the referee) gives a PD_3 -complex with 2-fold cover homotopy equivalent to $L(3, 1) \# -L(3, 1)$. (This group was first suggested as a test case in [2].)

1 A finite complex with group $S_3 *_{\mathbb{Z}/2\mathbb{Z}} S_3$

Let G be a group and let $\Gamma = \mathbb{Z}[G]$, $\varepsilon : C_1 = \Gamma \rightarrow \mathbb{Z}$ and $I(G) = \text{Ker}(\varepsilon)$ be the integral group ring, the augmentation homomorphism and the augmentation ideal, respectively. If M is a left Γ -module $\overline{M}$ shall denote the conjugate right module, with G -action given by $m.g = g^{-1}m$ for all $g \in G$ and $m \in M$, and similarly $\overline{N}$ shall denote the conjugate left module structure on a right Γ -module N . If C_* is a chain complex over Γ with an augmentation $\varepsilon : C_0 \rightarrow \mathbb{Z}$

a *diagonal approximation* is a chain homomorphism $\Delta : C_* \rightarrow C_* \otimes_{\mathbb{Z}} C_*$ (with diagonal G -action) such that $(\varepsilon \otimes 1)\Delta = id_{C_*} = (1 \otimes \varepsilon)\Delta$.

The cellular chain complex $C_*(\tilde{K})$ for the universal covering space of a finite 2-complex K determined by a presentation for a group is isomorphic to the Fox-Lyndon complex of the presentation, via an isomorphism carrying generators corresponding to based lifts of cells of K to the standard generators.

The symmetric group S_3 has a presentation $\langle a, b \mid a^2, abab^{-2} \rangle$. Let $\pi = S_3 *_{Z/2Z} S_3$, with presentation $\langle a, b, c \mid r, s, t \rangle$, where $r = a^2$, $s = abab^{-2}$ and $t = acac^{-2}$. The two obvious embeddings of S_3 into π admit retractions, as $\pi/\langle\langle b \rangle\rangle \cong \pi/\langle\langle c \rangle\rangle \cong S_3$. Let A , B and C be the cyclic subgroups generated by the images of a , b and c , respectively. The inclusions of A into S_3 and π induce isomorphisms on abelianization, while the commutator subgroups are $S'_3 = B$ and $\pi' = B * C$. Thus these groups are semidirect products: $S_3 \cong B \rtimes (Z/2Z)$ and $\pi \cong (B * C) \rtimes Z/2Z$. In particular, π is virtually free, and so has infinitely many ends. However it follows easily from the Grushko-Neumann Theorem that π is indecomposable. (See [3]).

The above presentations determine finite 2-complexes K and L , with fundamental groups S_3 and π , respectively. There are two obvious embeddings of K as a retract in L , with retractions $r_b, r_c : L \rightarrow K$ given by collapsing the pair of cells $\{c, t\}$ and $\{b, s\}$, respectively.

The chain complex $C_*(\tilde{K})$ has the form

$$\mathbb{Z}[S_3]^2 \xrightarrow{\partial_2} \mathbb{Z}[S_3]^2 \xrightarrow{\partial_1} \mathbb{Z}[S_3],$$

where $\partial_1(1, 0) = a - 1$, $\partial_1(0, 1) = b - 1$, $\partial_2(1, 0) = (a + 1, 0)$ and $\partial_2(0, 1) = (b^2a + 1, a - b - 1)$. The 2-chain $\psi = (a - 1, -ba + a + b^2 - b)$ is a 2-cycle, and so determines an element of $\pi_2(K) = H_2(\tilde{K}; \mathbb{Z})$, by the Hurewicz Theorem. Let $X = K \cup_{\psi} e^3$, and let C_* be the cellular chain complex for the universal cover $\tilde{X}$. (Thus $C_i = C_i(\tilde{K})$ for $i \leq 2$ and $C_3 \cong \mathbb{Z}[S_3]$). The dual cochain complex $C^* = Hom_{\Gamma}(C_*, \mathbb{Z}[S_3])$ is a complex of right $\mathbb{Z}[S_3]$ -modules.

We shall define new bases which display the structure of C_* to better advantage, as follows. Let $e_1 = (1, 0)$ and $e_2 = (-ba - b^2, 1)$ in C_1 and $f_1 = (1, 0)$ and $f_2 = (0, -a)$ in C_2 , and let g be the generator of C_3 corresponding to the top cell. Then $\partial_1 e_1 = a - 1$, $\partial_1 e_2 = -b^2a + ba + b^2 - 1$, $\partial_2 f_1 = (a + 1)e_1$, $\partial_2 f_2 = (b^2a + a - 1)e_2$, and $\partial_3 g = \psi = (a - 1)f_1 + (-b^2a + ba + b - 1)f_2$. The matrix for ∂_2 with respect to the bases $\{\tilde{e}_i\}$ and $\{\tilde{f}_j\}$ is diagonal, and is hermitian with respect to the canonical involution of $\mathbb{Z}[S_3]$, while the matrix for ∂_3 is the conjugate transpose for that of ∂_1 . Hence the chain complex $\overline{C^{3-*}}$

obtained by conjugating and reindexing the cochain complex C^* is isomorphic to C_* .

Let $\beta = b^2 + b + 1$ and $\nu = \sum_{s \in S_3} s = \beta(a + 1)$.

Lemma 1 *The complex X is a PD_3 -complex with $\tilde{X} \simeq S^3$.*

Proof Since C_* is the cellular chain complex of a 1-connected cell complex $H_0(C_*) \cong \mathbb{Z}$ and $H_1(C_*) = 0$. If $\partial_2(rf_1 + sf_2) = 0$ then $r(a + 1) = 0$ and $s(b^2a + a - 1) = 0$. Now the left annihilator ideals of $a + 1$ and $b^2a + a - 1$ in $\mathbb{Z}[S_3]$ are principal left ideals, generated by $a - 1$ and $(b - 1)(ba - 1)$, respectively. Hence $r = p(a - 1)$ and $s = q(b - 1)(ba - 1)$ for some $p, q \in \mathbb{Z}[B]$. A simple calculation gives $\partial_3((p(ba + b + 1) + q(ba + b))g) = rf_1 + sf_2$ and so $H_2(C_*) = 0$.

If $\partial_3hg = 0$ then $h(a - 1) = 0$, so $h = h_1(a + 1)$ for some $h_1 \in \mathbb{Z}[B]$, and $h(b^2a - ba - b + 1) = 0$. Now $h(b^2a - ba - b + 1) = h_1(1 - b)(a + b + 1)$, so $h_1(1 - b) = 0$. Therefore $h_1 = m\beta$ for some $m \in \mathbb{Z}$, so $h = m\nu$ and $H_3(C_*) = \mathbb{Z}[S_3]\nu g \cong \mathbb{Z}$. Hence $\tilde{X} \simeq S^3$. Now $H_3(X; \mathbb{Z}) = H_3(\mathbb{Z} \otimes_{\mathbb{Z}[S_3]} C_*) = \mathbb{Z}[1 \otimes g]$ and $tr([1 \otimes g]) = \nu g$, where $tr : H_3(X; \mathbb{Z}) \rightarrow H_3(\tilde{X}; \mathbb{Z})$ is the transfer homomorphism. The homomorphisms from $H^q(\tilde{C}^*)$ to $H_{3-q}(C_*)$ determined by cap product with $[X] = [1 \otimes g]$ may be identified with the Poincaré duality isomorphisms for $\tilde{X}$, and so X is a PD_3 -complex. $\square$

The verification that $\tilde{X} \simeq S^3$ is essentially due to [4] and the fact that X is a PD_3 -complex is due to [6]. The only novelty here is the diagonalization of ∂_2 , which was a guiding feature in the study of $\pi = S_3 *_{Z/2Z} S_3$.

Let $\Pi = \mathbb{Z}[\pi]$. The cellular chain complex for the universal covering space $\tilde{L}$ has the form

$$\Pi^3 \xrightarrow{\partial_2} \Pi^3 \xrightarrow{\partial_1} \Pi.$$

The differentials are given by $\partial_1(1, 0, 0) = a - 1$, $\partial_1(0, 1, 0) = b - 1$ and $\partial_1(0, 0, 1) = c - 1$, $\partial_2(1, 0, 0) = (a + 1, 0, 0)$, $\partial_2(0, 1, 0) = (b^2a + 1, a - b - 1, 0)$ and $\partial_2(0, 0, 1) = (c^2a + 1, 0, a - c - 1)$. In particular, $H_2(\tilde{L}; \mathbb{Z}) = \text{Ker}(\partial_2)$.

Let $\theta = (a - 1, -ba + a + b^2 - b, -ca + a + c^2 - c)$. Then $\partial_2(\theta) = 0$, and so θ determines an element of $\pi_2(L) = H_2(\tilde{L}; \mathbb{Z})$, by the Hurewicz Theorem. Let $Y = L \cup_{\theta} e^3$ and let D_* be the cellular chain complex for the universal covering space $\tilde{Y}$.

Let	Then
$\tilde{e}_1 = (1, 0, 0)$	$\partial_1 \tilde{e}_1 = a - 1$
$\tilde{e}_2 = (-ba - b^2, 1, 0)$	$\partial_1 \tilde{e}_2 = ba - b^2a + b^2 - 1$
$\tilde{e}_3 = (-ca - c^2, 0, 1)$	$\partial_1 \tilde{e}_3 = ca - c^2a + c^2 - 1$
$\tilde{f}_1 = (1, 0, 0)$	$\partial_2 \tilde{f}_1 = (a + 1)\tilde{e}_1$
$\tilde{f}_2 = (0, -a, 0)$	$\partial_2 \tilde{f}_2 = (b^2a + a - 1)\tilde{e}_2$
$\tilde{f}_3 = (0, 0, -a)$	$\partial_2 \tilde{f}_3 = (c^2a + a - 1)\tilde{e}_3$

Moreover $\theta = (a - 1)\tilde{f}_1 + (-b^2a + ba + b - 1)\tilde{f}_2 + (-c^2a + ca + c - 1)\tilde{f}_3$. Let $D^* = \text{Hom}_\Gamma(D_*, \Pi)$ be the cochain complex dual to D_* . Then it is easily seen that $\overline{D}^* \cong D_{3-*}$.

Theorem 2 *The complex Y is a PD_3 -complex.*

Proof Clearly $H_0(D_*) \cong \mathbb{Z}$ and $H_1(D_*) = 0$. The argument of the first part of Lemma 1 extends immediately to show that the kernel of ∂_2 is generated by $(a - 1)\tilde{f}_1$, $(b - 1)(ba - 1)\tilde{f}_2$ and $(c - 1)(ca - 1)\tilde{f}_3$. Hence these elements represent generators for $H_2(D_*)$. Let $\tilde{g}$ be the generator for D_3 corresponding to the top cell, so that $\partial_3 \tilde{g} = \theta$. Note that the image of g in $\mathbb{Z} \otimes_\varepsilon D_3$ is a cycle, and represents a generator for $H_3(Y; \mathbb{Z}) = H_3(\mathbb{Z} \otimes_\varepsilon D_*)$. If $h\theta = 0$ then (as in Lemma 1) $h = h_1(a + 1)$ for some $h_1 \in \mathbb{Z}[B * C]$ such that $h_1(b - 1) = h_1(c - 1) = 0$. It follows that $h_1 = 0$. Hence ∂_3 is injective and so $H_3(D_*) = 0$.

Let $\hat{1}$, $\hat{e}^*$, $\hat{f}^*$ and $\hat{g}$ denote the bases of D^* dual to the above bases for D_* . Let Δ be a diagonal approximation for D_* and suppose that $\Delta(\tilde{g}) = \sum_{0 \leq q \leq 3} \sum_{i \in I(q)} x_i \otimes y_i$, where $x_i \in D_q$ and $y_i \in D_{3-q}$, for all $i \in I(q)$ and $0 \leq q \leq 3$. Then $\sum_{i \in I(3)} x_i = \tilde{g}$. Let $r_i = \hat{g}(x_i)$ for $i \in I(3)$ and let $\tilde{\xi}$ denote the image of $\tilde{g}$ in $H_3(Y; \mathbb{Z}) = \mathbb{Z} \otimes_\varepsilon D_3$. Then $\varepsilon(\hat{g} \cap \tilde{\xi}) = \varepsilon(\sum_{i \in I(3)} \overline{r_i} y_i) = \varepsilon(\sum_{i \in I(3)} \overline{r_i}) = \varepsilon(\overline{\hat{g}(\tilde{g})}) = 1$, and so $\hat{g} \cap \tilde{\xi}$ generates $H_0(D_*)$. Since $H_1(D_*) = H_3(D_*) = H^0(\overline{D}^*) = H^2(\overline{D}^*) = 0$, $-\cap \tilde{\xi}$ induces isomorphisms $H^q(\overline{D}^*) \cong H_{3-q}(D_*)$ for all $q \neq 1$. The remaining case follows as in [5] from the facts that $\overline{D}^* \cong D_{3-*}$ and Δ is chain homotopic to $\tau\Delta$, where $\tau : D_* \otimes D_* \rightarrow D_* \otimes D_*$ is the transposition defined by $\tau(\alpha \otimes \omega) = (-1)^{pq} \omega \otimes \alpha$ for all $\alpha \in D_p$ and $\omega \in D_q$. Thus Y is a PD_3 -complex. $\square$

Can the last step of this argument be made more explicit? The work of Handel [1] on diagonal approximations for dihedral groups may be adapted to give the following formulae for a diagonal approximation for the truncation to degrees ≤ 2 of D_* which is compatible with the above two embeddings of K as a retract in L :

$$\begin{aligned}
\Delta(1) &= 1 \otimes 1 \\
\Delta(\tilde{e}_1) &= \tilde{e}_1 \otimes a + 1 \otimes \tilde{e}_1, \\
\Delta(\tilde{e}_2) &= \tilde{e}_2 \otimes 1 - ba\tilde{e}_1 \otimes (b-1) - b^2\tilde{e}_1 \otimes (b^2a-1) - (ba-b) \otimes ba\tilde{e}_1 \\
&\quad - (b^2-b) \otimes b^2\tilde{e}_1 + b \otimes \tilde{e}_2, \\
\Delta(\tilde{e}_3) &= \tilde{e}_3 \otimes 1 - ca\tilde{e}_1 \otimes (c-1) - c^2\tilde{e}_1 \otimes (c^2a-1) - (ca-c) \otimes ca\tilde{e}_1 \\
&\quad - (c^2-c) \otimes c^2\tilde{e}_1 + c \otimes \tilde{e}_3, \\
\Delta(\tilde{f}_1) &= \tilde{f}_1 \otimes 1 + \tilde{e}_1 \otimes a\tilde{e}_1 + 1 \otimes \tilde{f}_1, \\
\Delta(\tilde{f}_2) &= \tilde{f}_2 \otimes a + (b^2+b)\tilde{f}_1 \otimes (a-ba) + (b^2a+b^2)\tilde{f}_2 \otimes (a-ba) \\
&\quad + ((ba+b^2-1)\tilde{e}_1 + \tilde{e}_2) \otimes ((b^2a)\tilde{e}_1 + ba\tilde{e}_2) \\
&\quad - ((b^2a+1)\tilde{e}_1 + ba\tilde{e}_2) \otimes ((ba+a+b^2+b)\tilde{e}_1 + (b^2a+a)\tilde{e}_2) \\
&\quad - ((a+b)\tilde{e}_1 + b^2a\tilde{e}_2) \otimes ((ba+b^2)\tilde{e}_1 + a\tilde{e}_2) - (a+1)\tilde{e}_1 \otimes \tilde{e}_1 \\
&\quad + (a-b) \otimes (b^2+b)\tilde{f}_1 + (a-b) \otimes (b^2a+b^2)\tilde{f}_2 + a \otimes \tilde{f}_2 \quad \text{and} \\
\Delta(\tilde{f}_3) &= \tilde{f}_3 \otimes a + (c^2+c)\tilde{f}_1 \otimes (a-ca) + (c^2a+c^2)\tilde{f}_3 \otimes (a-ca) \\
&\quad + ((ca+c^2-1)\tilde{e}_1 + \tilde{e}_3) \otimes ((c^2a)\tilde{e}_1 + ca\tilde{e}_3) \\
&\quad - ((c^2a+1)\tilde{e}_1 + ca\tilde{e}_3) \otimes ((ca+a+c^2+c)\tilde{e}_1 + (c^2a+a)\tilde{e}_3) \\
&\quad - ((a+c)\tilde{e}_1 + c^2a\tilde{e}_3) \otimes ((ca+c^2)\tilde{e}_1 + a\tilde{e}_3) - (a+1)\tilde{e}_1 \otimes \tilde{e}_1 \\
&\quad + (a-c) \otimes (c^2+c)\tilde{f}_1 + (a-c) \otimes (c^2a+c^2)\tilde{f}_3 + a \otimes \tilde{f}_3
\end{aligned}$$

These formulae were derived from the work of Handel by using the canonical involution of $\mathbb{Z}[S_3]$ to switch right and left module structures and showing that C_* is a direct summand of a truncation of the Wall-Hamada resolution for S_3 . (In Handel's notation $a = y$, $b = x$, $e_1 = c_1^2$, $e_2 = -c_1^1 - c_1^2(x + xy)$, $f_1 = c_2^3$, $f_2 = -c_2^1y + c_2^2x^2 - c_2^3y$ and $g = -(c_3^1 + c_3^3)(x + y) - c_3^4y$). Handel's work also leads to a formula for $\Delta(g)$, but it is not clear what $\Delta(\tilde{g})$ should be.

2 Other PD_3 -complexes with this group

Having constructed one PD_3 -complex with group π one may ask how many there are. Any such PD_3 -complex must be orientable. For let $w_1 : \pi \rightarrow \{\pm 1\}$ be a homomorphism and define an involution on Γ by $\bar{g} = w_1(g)g^{-1}$, for all $g \in \pi$. Let $w = w_1(a)$ and $R = \mathbb{Z}[\pi/\pi'] = \mathbb{Z}[a]/(a^2 - 1)$. Let $J = \text{Coker}(\overline{\partial}_2^{tr})$,

where $\partial_2 : \Pi^3 \rightarrow \Pi^3$ is the presentation matrix for $I(\pi)$ given in §1. Then $R \otimes_{\Gamma} I(\pi) \cong R/(a+1) \oplus (R/(a+1, 3))^2$, while $R \otimes_{\Gamma} J \cong R/(a+w) \oplus (R/(a+w, 3))^2$. If the pair (π, w_1) is realized by a PD_3 -complex then $I(\pi)$ and J are projective homotopy equivalent [5]. But then $R \otimes_{\Gamma} I(\pi)$ and $R \otimes_{\Gamma} J$ are projective homotopy equivalent R -modules, and so we must have $w = 1$.

If W is an oriented PD_3 -complex with fundamental group G and $c_W : W \rightarrow K(G, 1)$ is a classifying map let $\mu(W) = c_{W*}[W] \in H_3(W; \mathbb{Z})$. Two such PD_3 -complexes W_1 and W_2 are homotopy equivalent if and only if $\mu(W_1)$ and $\mu(W_2)$ agree up to sign and the action of $Out(G)$. Turaev constructed an isomorphism ν_C from $H_3(G; \mathbb{Z})$ to a group $[F^2(C), I(G)]$ of projective homotopy classes of module homomorphisms and showed that $\mu \in H_3(G; \mathbb{Z})$ is the image of the orientation class of a PD_3 -complex if and only if $\nu_C(\mu)$ is the class of a self-homotopy equivalence [5].

When $G = \pi = S_3 *_{Z/2Z} S_3$ we have $F^2(C) \cong I(\pi)$, and $H_3(\pi; \mathbb{Z}) \cong H_3(\pi'; \mathbb{Z}) \oplus H_3(Z/2Z; \mathbb{Z}) \cong (Z/3Z)^2 \oplus (Z/2Z)$. Let W' be the double cover of W , with fundamental group $\pi' \cong (Z/3Z) * (Z/3Z)$. Then W' is a connected sum, by Theorem 1 of [5], and so it is homotopy equivalent to one of the 3-manifolds $L(3, 1) \# L(3, 1)$ and $L(3, 1) \# -L(3, 1)$. (These may be distinguished by the torsion linking forms on their first homology groups). In particular, $\mu(W')$ has nonzero entries in each summand. Since $\mu(W')$ is the image of $\mu(W)$ under the transfer to $H_3(\pi'; \mathbb{Z}) \cong (Z/3Z)^2$ the image of $\mu(W)$ in each $Z/3Z$ -summand must be nonzero. Let $u \in H^1(W; \mathbb{F}_2)$ correspond to the abelianization homomorphism. Since $\beta_2(W; \mathbb{F}_2) = \beta_1(W; \mathbb{F}_2) = 1 = \beta_2(\pi; \mathbb{F}_2)$ we have $u^2 \neq 0$, and so $u^3 \neq 0$, by Poincaré duality. It follows easily that the image of $\mu(W)$ in the $Z/2Z$ -summand must be nonzero also. (Note that W' is $\mathbb{Z}_{(2)}$ -homology equivalent to S^3 and so W is $\mathbb{Z}_{(2)}[Z/2Z]$ -homology equivalent to RP^3). Since reversing the orientation of W reverses that of W' , we may conclude that there are at most two distinct homotopy types of PD_3 -complexes with fundamental group π , and that they may be detected by their double covers.

The retractions r_b and r_c of L onto K extend to maps $r_b, r_c : Y \rightarrow X$. These maps induce the same isomorphism $H_3(Y; \mathbb{Z}) \cong H_3(X; \mathbb{Z})$, and so their lifts to the double covers induce the same isomorphism $H_3(Y'; \mathbb{Z}) \rightarrow H_3(X'; \mathbb{Z})$. Hence $Y' \simeq L(3, 1) \# L(3, 1)$, rather than $L(3, 1) \# -L(3, 1)$. The referee has pointed out that if we use $\xi = (a-1)\tilde{f}_1 + (-b^2a + ba + b - 1)\tilde{f}_2 - (-c^2a + ca + c - 1)\tilde{f}_3$ instead of θ (changing only the sign of the final term) then $Z = L \cup_{\xi} e^3$ is another PD_3 -complex with $\pi_1(Z) \cong \pi$, and a similar argument shows that the double cover is now $Z' \simeq L(3, 1) \# -L(3, 1)$.

The question of whether every aspherical PD_3 -complex is homotopy equivalent to a 3-manifold remains open. The recent article [7] gives a comprehensive

survey of Poincaré duality in dimension 3, emphasizing the role of the JSJ decomposition in relation to this question.

Acknowledgement I would like to thank the referee for suggesting the modification giving the example covered by $L(3, 1) \sharp -L(3, 1)$, and for other improvements to the exposition.

References

- [1] Handel, D. On products in the cohomology of the dihedral groups, Tôhoku Math. J. 45 (1993), 13-42. MR1200878
- [2] Hillman, J.A. On 3-dimensional Poincaré duality complexes and 2-knot groups, Math. Proc. Cambridge Phil. Soc. 114 (1993), 215-218. MR1230128
- [3] Hillman, J.A. An indecomposable PD_3 -complex whose group has infinitely many ends, Math. Proc. Cambridge Phil. Soc., to appear (2005).
- [4] Swan, R.G. Periodic resolutions for finite groups. Ann. of Math. 72 (1960), 267-291. MR0124895
- [5] Turaev, V.G. Three-dimensional Poincaré complexes: homotopy classification and splitting. Math. USSR-Sb. 67 (1990), 261-282. MR1015042 (91c:57031) Turaev, V. G. Three-dimensional Poincaré complexes: homotopy classification and splitting. (Russian) Mat. Sb. 180 (1989), no. 6, 809–830.
- [6] Wall, C.T.C. Poincaré complexes: I. Ann. of Math. 86 (1967), 213-245. MR0217791
- [7] Wall, C.T.C. Poincaré duality in dimension 3, in *Proceedings of the Casson Fest (Arkansas and Texas 2003)* (edited by C.McA.Gordon and Y.Rieck), Geom. Topol. Monogr. 7 (2004), 1-26.

*School of Mathematics and Statistics F07
University of Sydney, NSW 2006, Australia*

Email: `johnh@maths.usyd.edu.au`

Received: 4 August 2004